November 23 – Dec. 1, 2013 Kentucky Waterfowl Ground Counts

Overall, wetland conditions are much improved from last year at this time. Many of the public waterfowl areas report moist soil units as being either at seasonal norms. While agricultural crops were impacted by the spring and summer rains, natural vegetation was very robust and produce and excellent source of natural foods that should persists well into winter meeting the seasonal demands of migrating and wintering waterfowl. An early cold front should provide a strong early push of mallards that have been held up in the central Illinois and Missouri River bottoms over the last week to 10 days. The influx of migrating birds coupled with good to excellent wetland conditions should provide outstanding conditions for the season opener.

The following list below, represent the results of ground counts made by local KDFWR wildlife biologists. The locations and counts presented below probably represent a minimal number of what can be expected listed on the surveyed areas. Good hunting!

| Location | Number | Species Composition | Comments |
|------------------------|----------------|--------------------------------|-------------------------|
| Ballard WMA | 20,290 ducks | 85% mallard, 10% gadwall, | MSUs partially filled. |
| | 90 CG geese | remainder a mix of wigeon, | Main Ohio River pump |
| | 990 W-F geese | pintail, green-winged teal and | on and hunt units are |
| | 450 Snow geese | wood duck. | filling. |
| Barren River Lake | 300 ducks | 80% divers, diverse species | Lake 4.9' above winter |
| | | composition. Ring-necked | pool. Water level |
| | | ducks, redhead and ruddy | dropping |
| | | most common | |
| Birdsville Island | No count | | |
| Kentucky/Barkley Lakes | No Count | | |
| Duck Island | 2,000-3,000 | 85% mallard, 10% green- | No row crops, excellent |
| | ducks | winged teal, and 5% a mix of | moist soil. Water and |
| | | gadwall, pintail and shoveler | habitat good/excellent |
| Boatwright WMA | 1,576 ducks | 96% mallards remainder wood | MSU's have good water |
| | | ducks, gadwall and others | on hunt units. Several |
| | | Canada geese | ducks observed moving |
| | | | around. |
| Clear Creek WMA | No Count | | |
| Peabody WMA | 385 ducks | 80% mallard, 15% other | Plenty of water, MSU's |
| | 50 coots | dabblers | near capacity. Good |
| | 236 Can. Geese | | stand of natural veg. |
| Sloughs WMA | 3,500 ducks | 85% mallards remainder | Most birds were on the |
| | 2,500 WFG | wigeon, gadwall, black duck. | refuge, feeding on |
| | 2,500 SG | | winter wheat. Wells |
| | 250 CG | | running, 85% of MSU's |
| | 17 Swan | | at capacity. Natural |
| | | | areas look good. |

| Yellowbank WMA | 570 | Mostly mallards and wood | Water level at |
|-----------------------|-----------|-------------------------------|-------------------------|
| | | ducks Some greeni-winged teal | maximum capacity on |
| | | and gadwall | most MSU's. |
| Blue Grass Army Depot | 110 ducks | 50% mallard, 20% gadwall, | All water at seasonal |
| (all lakes and ponds) | | 15% black duck, remainder | levels. Lake Gen about |
| | | divers | 2/3 full |
| Herrington Lake | 0 | No ducks observed on lake or | |
| | | ash pond. Some mallards | |
| | | observed on local farm ponds | |
| Taylorsville Lake | 50 CG | Canada geese observed on | Lake above winter pool |
| | | adjacent farm pond. | and falling. Pumping |
| | | | underway on MSUs. |
| Cedar Cr./Dix R. WMA | 10 ducks | Mallard | Wetland units on Dix |
| | 500 coots | | River WMA are partially |
| | | | flooded. |
| Green River WMA | 200 ducks | 90% mallards with some GWT | All MSU's and beaver |
| | | and wigeon. | ponds full. Lake above |
| | | | winter pool |
| Cave Run Lake | 400 | Mostly gadwall, some mallard, | Zilpo Flats and Dam |
| | | wigeon and pintail. | area. |
| | 100 | Unknown duck | |
| | 125 | Canada geese | |
| | 800 | Coots | |